



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$3a^2(m - \frac{1}{m}) + 4ay + (x^2 + y^2 - 4ax)(m - \frac{1}{m}) = 0; \text{ or}$$

$$3a^2p + 4ay + (x^2 + y^2 - 4ax)p = 0 \dots (II).$$

From (II), we get  $p = -\frac{4ay}{x^2 + y^2 - 4ax + 3a^2}$ , and substituting this in (I), we get

$$(x^2 + y^2 + 3a^2 - 4ax)^2 (3a^2 + 10ax - x^2 - y^2) - 16a^2y^2(x^2 + y^2 - 4ax) = 0.$$

Also solved by M. A. Harding, S. Lefschetz, and A. H. Holmes.

367. Proposed by E. B. ESCOTT, University of Michigan, Ann Arbor, Michigan.

Solve the simultaneous equations:

$$\frac{2x}{1+x^2} = y \dots (1); \quad \frac{2y}{1+y^2} = z \dots (2); \quad \frac{2z}{1+z^2} = u \dots (3); \quad \frac{2u}{1+u^2} = x \dots (4).$$

Solution by PROFESSOR F. L. GRIFFIN, Reed College, Portland, Oregon.

By inspection three solutions are  $x=y=z=u=1$ , or  $-1$ , or  $0$ ; and there can be but fourteen others. Now let  $x=i \tan \theta$ , [ $i=\sqrt{-1}$ ], whence  $y=i \tan 2\theta$ ,  $z=i \tan 4\theta$ ,  $u=i \tan 8\theta$ , and  $x=i \tan 16\theta$ . But  $\tan 16\theta = \tan \theta$  for finite values of  $\theta$  only if  $16\theta = \theta + n\pi$ , or  $\theta = n\pi/15$ . Thus we have

$$\begin{array}{llllll} x=0, & i \tan \pi/15, & i \tan 2\pi/15, & i \tan 3\pi/15, & \dots, & i \tan 14\pi/15; \\ y=0, & i \tan 2\pi/15, & i \tan 4\pi/15, & i \tan 6\pi/15, & \dots, & i \tan 28\pi/15; \\ z=0, & i \tan 4\pi/15, & i \tan 8\pi/15, & i \tan 12\pi/15, & \dots, & i \tan 56\pi/15; \\ u=0, & i \tan 8\pi/15, & i \tan 16\pi/15, & . & . & . \end{array}$$

the values for  $y, z, u$  being of course the same sets as for  $x$  in different orders. The values  $x=+1, -1$  correspond to infinite values of  $\theta$  for which  $\tan \theta = -i, +i$ .

## GEOMETRY.

393. Proposed by S. LEFSCHETZ, University of Nebraska.

Draw a triangle having a given angle, and with its vertices on three given concentric circles.

I. Solution by H. C. FEEMSTER, A. B., Professor of Mathematics, York College, York, Nebraska.

Let  $x^2 + y^2 = c^2$ ,  $x^2 + y^2 = b^2$ ,  $x^2 + y^2 = a^2$ , be the given circles, and  $\phi$  the given angle. Place the vertex of the given angle,  $\phi$ , on the circumference of the first circle at  $O$ ,  $c$ , the angle being formed by the lines,